

# Conservación Colombiana

Número 13 • 18 Mayo 2010

©2010 Fundación ProAves • Bogotá • Colombia • ISSN 1900-1592



**A new species of Antpitta from the Colibrí del Sol  
Bird Reserve, Colombia /**

**Revision of the status of bird species in Colombia**

---

**Una nueva especie de tororoi de la Reserva Natural  
de las Aves Colibrí del Sol, Colombia /**  
**Revisión del estatus de especies de aves en Colombia**

## **Conservación Colombiana**

Revista de difusión de las acciones de conservación de la biodiversidad en Colombia.

ISSN 1900–1592

Entidad sin ánimo de lucro S0022872 – Cámara de Comercio de Bogotá

**Conservación Colombiana** es una revista científica publicada por la Fundación ProAves, institución que tiene como misión «proteger las aves silvestres y sus hábitat en Colombia a través de la investigación, acciones de conservación puntuales y el acercamiento a la comunidad». El propósito de la revista es divulgar las acciones de conservación que se llevan a cabo en Colombia, para avanzar en su conocimiento y en las técnicas correspondientes. El formato y tipo de los manuscritos que se publican es variado, incluyendo reportes de las actividades de conservación desarrolladas, resultados de las investigaciones y el monitoreo de especies amenazadas, proyectos de grado de estudiantes universitarios, inventarios y conteos poblacionales, planes de acción o estrategias desarrolladas para especies particulares, sitios o regiones y avances en la expansión de la red de áreas protegidas en Colombia. Conservación Colombiana está dirigida a un público amplio, incluyendo científicos, conservacionistas y personas en general interesadas en la conservación de las especies amenazadas de Colombia y sus hábitats.

Fundación ProAves – Colombia

Dirección: Carrera 20 No. 36–61, La Soledad, Bogotá

Teléfonos: (1) 245 5134 – 340 3239

Fax: (1) 340 3285

[www.ProAves.org](http://www.ProAves.org)

## **Comité Editorial**

David Caro, Lina Daza, Thomas Donegan, Alonso Quevedo y Paul Salaman

## **Fotografía portada**

Holotype Fenwick's Antpitta – *Grallaria fenwickorum* sp nov. Reserva Natural de Aves Colibrí del Sol Urrao, Antioquia, Colombia © Fundación ProAves [www.proaves.org](http://www.proaves.org).

## **Permisos y derechos de autor**

Toda reproducción parcial o total de esta obra está prohibida sin el permiso escrito de los autores y de la Fundación ProAves. Publicado el 18 Mayo 2010 en papel y en línea.

**Conservación Colombiana** está cobijada por la ley colombiana de derechos de autor, Ley 23 de 1982, Ley 44 de 1993 y Decisión 351 de la Comisión del Acuerdo de Cartagena de 1993.

**Conservación Colombiana** es publicada gracias a la Ley para la Conservación de Especies Migratorias / *Neotropical Migratory Bird Conservation Act* (NMBCA) del *US Fish and Wildlife Service* y American Bird Conservancy.



## Contenidos. Contents.

<b>Conservación Colombiana 13</b>	
<b>Editorial sobre la descripción de una nueva especie de <i>Grallaria</i>.</b> Editorial about the description of a new species of <i>Grallaria</i> . <i>Comité Editorial de Conservación Colombiana.</i>	<b>4</b>
<b>A new species of Antpitta (family Grallariidae) from the Colibrí del Sol Bird Reserve, Colombia.</b> Una nueva especie de tororoi (familia Grallariidae) de la Reserva Natural de Aves Colibrí del Sol, Colombia. <i>Luis Felipe Barrera, Avery Bartels, &amp; Fundación ProAves de Colombia</i>	<b>8</b>
<b>Revision of the status of bird species occurring or reported in Colombia 2010.</b> Revisión del estatus de especies de aves que ocurren o son reportados en Colombia 2010. <i>Thomas Donegan, Paul Salaman, David Caro &amp; Miles McMullan</i>	<b>25</b>
<b>Artículos publicados anteriormente únicamente en línea, imprimidos en esta edición. Articles previously published only online, printed in this edition.</b>	
<b>Conservación Colombiana 5</b>	
<b>Sixteen bird species new for Colombia.</b> Dieciséis especies de aves nuevas para Colombia. <i>Paul Salaman, Nick Bayly, Robert Burridge, Mark Grantham, Mark Gurney, Alonso Quevedo, Luis Eduardo Urueña &amp; Thomas Donegan</i> <i>Conservación Colombiana 5: 80-85</i>	<b>5: 80</b>
<b>Conservación Colombiana 8</b>	
<b>Revision of the status of various bird species occurring or reported in Colombia.</b> Revisión del estatus de especies de aves que ocurren o son reportados en Colombia. <i>Thomas Donegan, Paul Salaman &amp; David Caro</i> <i>Conservación Colombiana 8: 80-86</i>	<b>8: 80</b>
<b>First confirmed locality of Yellow-throated Warbler <i>Dendroica dominica</i> for Colombia.</b> Primer localidad confirmada para <i>Dendroica dominica</i> en Colombia. <i>Trevor Ellery, Deb Hahn, Guy Foulks, Christian Olaciregui &amp; Thomas Donegan</i> <i>Conservación Colombiana 8: 87</i>	<b>8: 87</b>
<b>Conservación Colombiana 13: Suplemento</b>	
<b>Listado de las aves de Colombia 2010.</b> Checklist of the birds of Colombia 2010. <i>Paul Salaman, Thomas Donegan &amp; David Caro</i>	<b>online</b>

## Editorial sobre la descripción de una nueva especie de *Grallaria*

---

El presente editorial tiene como objetivo explicar la historia del artículo en esta revista en el cual se describe una nueva especie de la familia Grallariidae *Grallaria* para la ciencia.

Durante una exploración apoyada por la Fundación ProAves de Colombia en Agosto del 2004 en el páramo del sol, Cordillera Occidental, Urrao, Antioquia, se realizaron importantes y significativos descubrimientos para la avifauna Colombiana, dentro de los cuales tenemos: el redescubrimiento de una especie de colibrí “perdido” (*Coeligena orina*) (Krabbe et al. 2005), los primeros datos sobre la voz del tapaculo de paramillo (*Scytalopus canus*) (Krabbe & Cadena 2010) y una nueva población del loro amenazado *Hapalopsittaca amazonina* (Krabbe et al. 2006). Los registros más importantes de estas exploraciones iniciales han sido publicados (Flórez et al. 2004, Krabbe et al. 2006). Entre estos resultados, Krabbe et al. (2006) también reportaron varias subespecies de aves que podían ser nuevas para la ciencia.

Debido a la gran cantidad de especies de aves de importancia para la conservación y la presencia de posibles nuevas aves en esta zona, la Fundación ProAves decidió establecer una reserva natural en el páramo del sol. Las tierras fueron compradas con fines de conservación a perpetuidad en el año 2005 y designadas como una reserva natural de las aves de la Fundación ProAves con su nombre “RNA Colibrí del Sol”. Con el ánimo de consolidar los procesos de conservación en esta importante reserva, el 1 de febrero del año 2008, la Fundación ProAves contrató al biólogo Diego Andrés Carantón, como coordinador de la reserva natural de las aves Colibrí del Sol para manejarla y realizar estudios adicionales en la misma.

Teniendo en cuenta lo anterior, es preciso traer en cuestión algunas de las cláusulas y condiciones

que se pactaron entre la Fundación ProAves de Colombia y el señor Carantón dentro del contrato de trabajo celebrado, las cuales incluyen las siguientes cláusulas:

A) En la cláusula primera del contrato se indicó que: *LA EMPLEADORA contrata los servicios personales del TRABAJADOR(A) y este se obliga a: b) A no prestar directa ni indirectamente servicios laborales a otros EMPLEADORES, ni a trabajar por cuenta propia en el mismo oficio, durante la vigencia de este contrato. Observando en su trabajo el cuidado y la diligencia necesarios para el correcto desempeño de su labor en forma personal y directa.*

B) En la cláusula Cuarta del contrato arriba mencionado, se dispuso que el trabajador debe: “*apoyar la prórroga y el cumplimiento de las obligaciones del permiso de investigación de la Corporación Autónoma Regional en la Zona en coordinación con el encargado de convenios y Representante Legal*”, “*presentar un reporte mensual entre el 25 y el 30 de cada mes*” al director de áreas protegidas, dirección ejecutiva, director de investigación y coordinador de base de datos de la Fundación. Del mismo modo y en relación a las obligaciones en el área de investigación, el trabajador como investigador se comprometió a realizar “*Anillamiento. Realizar capturas, anillar y tomar los datos establecidos en el protocolo metodológico con 15 redes de niebla en tres estaciones de monitoreo seleccionadas por tipo de vegetación y/o rango de elevación durante seis (6) días (2 días x cada estación) para un total de 540 horas /red al mes. Importante ubicar dos series de redes por estación de monitoreo para generar un cambio cada mes con el fin de evitar que las aves se acostumbren a la presencia de redes. De ser necesaria la colección debido a muertes accidentales los individuos colectados serán preparados adecuadamente, debidamente*

*etiquetados y reportados a la coordinadora del programa para el respectivo procedimiento de traslado a la respectiva colección ornitológica.... Fotografía:. Llevar un registro fotográfico de las especies capturadas, procurando mantener representatividad en cuanto a registros de juveniles, machos y hembras cuando sea posible el cual debe ser enviado mensualmente vía correo físico a la Fundación en un disco compacto. Envío Datos. Archivar correctamente los datos originados durante el monitoreo y enviar fotocopias legibles por correo identificando a la oficina de la fundación en Bogotá para su digitación. Los formularios de datos tomados a la fecha deben llegar entre el 25 del mes que finaliza y los cinco primeros días del mes siguiente. Formularios con letra legible o malas fotocopias no serán aceptados. ... Entregar personalmente los datos en sus formatos originales en la oficina de ProAves o personalmente a cualquier miembro del equipo que viaje a Bogotá”.*

C) En la cláusula Tercera se determinó que el trabajador estaba obligado a: “*observar rigurosamente las normas que le fije La Fundación*”, “*cumplir estrictamente las instrucciones que le sean dadas por La Fundación, o por quienes la representan, respecto del desarrollo de sus actividades*” y “*cumplir permanentemente con el espíritu de lealtad, colaboración y disciplina con La Fundación*”.

D) En la cláusula décima séptima del contrato se dispuso que: “*Las obras, invenciones y/o descubrimientos realizados por EL TRABAJADOR(A) durante o con ocasión de la ejecución del presente Contrato, pertenecen a LA EMPLEADORA, de conformidad con el artículo 539 del Código de Comercio, así como el artículo 20 y concordantes de la Ley 23 de 1982 sobre derechos de autor*”.

E) En la cláusula Décima Tercera se indicó que: “*Este contrato ha sido redactado estrictamente de acuerdo con la ley y la jurisprudencia y será interpretado de buena fe y en consonancia con el Código Sustantivo del*

*Trabajo cuyo objeto, definido en su artículo 1, es lograr la justicia en las relaciones entre EMPLEADORES Y TRABAJADORES dentro de un espíritu de coordinación económica y equilibrio social”.*

Es claro en este punto que el señor Diego Carantón como trabajador de ProAves, tenía una serie de compromisos y obligaciones que acatar y que las actividades que realizó como investigador estaban sujetas al vínculo como trabajador de ProAves. Del mismo modo es preciso señalar que la Fundación cumplió con todas sus obligaciones de acuerdo a la Ley y explícitas en el contrato, fue así como el señor Carantón recibió de manera puntual su salario, control y pago mensual de los debidos aportes a la seguridad Social, pago de prestaciones sociales y otros beneficios como empleado de ProAves.

En la segunda semana del mes de Octubre del año 2008 la Fundación se enteró, lamentablemente por terceros, del hecho de que Diego Carantón había colectado (sacrificado) dos individuos de una nueva especie de *Grallaria*, con propósitos de utilizar los especímenes como base de una descripción. El mismo hecho, infiere una violación de las estipulaciones pactadas en el contrato suscrito con la Fundación, ya que el señor Carantón no informó realmente a la fundación de las actividades que venía realizando, en razón a que el mismo no incluyó datos sobre la nueva especie en sus reportes mensuales, tampoco realizó coordinación con la fundación relacionada con la colecta o el depósito de especímenes. Además, el investigador había desarrollado planes de tratar la información producto de sus investigaciones de forma independiente. El proceso ocurrió sin tener en cuenta los derechos sobre la propiedad intelectual de la Fundación explícitos en el contrato, teniendo en cuenta, que se destinó como apoyo otro empleado de la fundación (sin este conocer la falla del reglamento que se venia comentiendo), utilización de equipos de investigación de propiedad de la Fundación para

realizar esta acción, y el lugar de la investigación: el interior de un área de propiedad de la fundación destinada para la conservación a perpetuidad de todos sus organismos que allí habitan.

El descubrimiento de esta especie es motivo de orgullo para la Fundación *ya que fue el resultado de investigaciones financiadas por la Fundación así como que el hallazgo se realizó dentro de una reserva natural de la Fundación*. Tradicionalmente, las descripciones de nuevas especies cuentan con un “espécimen tipo” de la nueva especie, el cual se deposita en un museo, así que se entiende por qué el investigador se interesó en colectar dos especímenes de esta población con fines de realizar una descripción. Sin embargo, la colecta se realizó sin la autorización de la Fundación ProAves y sin el respectivo permiso de colecta ante la entidad competente, en este caso la Corporación Autónoma Regional de Urabá (CORPOURABA). Por otro lado, el investigador tampoco consiguió la necesaria “constancia de depósito” sobre los especímenes depositados en el Instituto de Ciencias Naturales (Universidad Nacional), ni lo reportó a la Fundación como obligación de acuerdo a los términos estipulados dentro de su contrato vigente durante esta acción. La colecta sin permiso de dichos especímenes, ha causado inconvenientes para las relaciones entre la Fundación y las corporaciones autónomas, por lo cual el reglamento interno de la Fundación exige dentro de sus procedimientos la tramitación de los debidos permisos antes de proceder con las actividades en el terreno y entrega de la constancia de depósito a la Fundación por su posterior depósito por parte de la Fundación con la corporación autónoma relevante. Actualmente CORPOURABA se encuentra pendiente de este proceso y situación, por lo que está realizando su propia investigación.

Diego Carantón cuando entregó el informe mensual a la Fundación en septiembre del 2008 (y en otros informes anteriores) no mencionó la

colecta de individuos de *Grallaria*, ni del posible descubrimiento de una nueva especie para la ciencia, a pesar de los términos y cláusulas de su contrato laboral. Igualmente, datos sobre las capturas no se encuentran registrados en los datos de anillamiento y tampoco se registraron fotos de la nueva especie en el material entregado a la Fundación. Sin embargo, el señor Carantón, con otros co-autores ajenos a la Fundación, prepararon un artículo científico sobre la descripción de la nueva especie. Después de haber sido notificado sobre la situación, miembros de la Fundación ProAves se acercaron al señor Carantón y, al parecer, estuvieron cerca de llegar a un acuerdo sobre la publicación de la nueva especie en conjunto y sobre el apropiado reconocimiento de la contribución de la Fundación a este hallazgo. No obstante, varias semanas después, el señor Carantón de forma radical cambió de opinión, suprimió del artículo los nombres de autores investigadores vinculados con la Fundación ProAves y agradecimientos a la Fundación por su apoyo a su investigación y envió un manuscrito de un artículo sobre la descripción de una nueva especie a la revista norteamericana *Condor*.

En diciembre del 2009, durante el proceso de “peer review” de un manuscrito del artículo sobre la descripción de una nueva especie a la revista norteamericana *Condor*, los editores de la revista se contactaron con ProAves para preguntar sobre el permiso de colecta. Cuando se había explicado la situación y irregularidades presentadas, la revista *Condor* rechazó el artículo hasta que la situación entre el autor y ProAves se resolviese. A pesar de que ProAves intentó conciliar con el señor Carantón durante nueve (9) oportunidades con el propósito de resolver la situación, desafortunadamente hasta el día de hoy no se ha podido llegar a ningún tipo de acuerdo con el señor Carantón.

La publicación del artículo sobre la descripción de la nueva especie en esta edición de *Conservación Colombiana*, no tiene como su

propósito desmeritar el hecho de que el biólogo Diego Carantón, con el apoyo del actual guardabosque de la Reserva Colibri del sol el señor Luis Rubelio García, fueron las personas quienes primero se percataron de que la población correspondía a una posible especie nueva de *Grallaria*. No obstante, el personal de ProAves vinculado con el asunto, nos han presentado historias relacionadas con las inconsistencias sobre el hallazgo por parte del señor Carantón detallados arriba. A pesar de que varias personas vinculadas con ProAves han intentado de una u otra forma mediar y llegar a un acuerdo, lastimosamente no se logró a pesar de la gravedad de la situación y teniendo en cuenta la infracción e incumplimiento cometida por el señor Caranton dentro del tiempo que estuvo contratado. Aunque ProAves en general incluye a todos sus investigadores relevantes como autores en las muchísimas publicaciones resultado de investigaciones apoyadas por la Fundación a través de los últimos 10 años, en el presente caso, lastimosamente, no ha sido factible incluir a Diego Carantón como autor. No obstante, cualquier derecho sobre propiedad intelectual relacionado con el descubrimiento de esta nueva especie, pertenece a la Fundación ProAves de Colombia y no existe ningún tipo de restricción o impedimento para utilizarlos en la descripción aquí publicada.

Los autores del artículo sobre la descripción de *Grallaria* publicado en esta edición de Conservación Colombiana han sometido un artículo, el cual ha sido sujeto a reseña de colegas y aceptado por su publicación por el comité editorial. Se espera que con esta publicación se

cierre esta incómoda situación. Es de aclarar que la publicación del nombre y detalles sobre la morfología, comportamiento y conservación de la nueva especie no perjudican la publicación de análisis científicos u otros datos sobre esta especie que puede ser realizado por otros autores en el futuro.

Así, con esta especie ya descrita, se espera fomentar más medidas de conservación para la nueva especie de *Grallaria* y su hábitat, teniendo en cuenta que las descripciones son frecuentemente un portal para medidas como la designación de especies amenazadas, zonas de importancia de conservación y fondos para la investigación y conservación.

### Comité Editorial, Conservación Colombiana

### Referencias

- Flórez, P., Krabbe, N., Castaño, J., Suárez, G. & Arango, J.D. 2004. Evaluación Avifauna del Páramo de Frontino, Antioquia, Agosto 2004. *Colombian EBA Project Report Series 6*.
- Krabbe, N. & Cadena, C. D. 2010. A taxonomic revision of the Paramo Tapaculo *Scytalopus canus* Chapman (Aves: Rhinocryptidae), with description of a new subspecies from Ecuador and Peru. *Zootaxa* 2354: 56–66.
- Krabbe, N., Flórez, P., Suárez, G., Castaño, J., Arango, J.D., Pulgarín, P.C., Múnera, W.A., Stiles, F.G., Salaman, P. 2005. Rediscovery of the Dusky Starfrontlet *Coeligena orina*, with a description of the adult plumages and a reassessment of its taxonomic status. *Ornitología Colombiana* 3:28-35
- Krabbe, N., Flórez, P., Suárez, G., Castaño, J., Arango, J. D. & Duque, A. 2006. The birds of Páramo Frontino, West Andes of Colombia. *Orn. Colombiana* 4: 39–50.

# A new species of Antpitta (family Grallariidae) from the Colibrí del Sol Bird Reserve, Colombia

Luis Felipe Barrera, Avery Bartels, & Fundación ProAves de Colombia

Fundación ProAves de Colombia, Cra 20 No. 36-61, Bogotá, Colombia. Email: fundacion@proaves.org

## Abstract

A new species of Antpitta *Grallaria* (family Grallariidae) is described from the Colibrí del Sol Bird Reserve situated on the southeastern slope of the Páramo del Sol massif on the northern end of the Western Andes of Colombia. Fenwick's Antpitta *Grallaria fenwickorum* sp nov. most closely resembles Brown-banded Antpitta *G. milleri*, which is endemic to the Central Andes of Colombia, but it has a slate grey breast, lacking the brown breast band and flanks of *G. milleri*. The song of the new species involves three notes of increasing length and acoustic frequency. The individual notes are shorter, lower and of a different note shape compared to *G. milleri*. In addition, the new species' call is shorter and lower pitched. Fenwick's Antpitta is restricted to upper montane cloud forest (2,600-3,300 m) where territories are largely restricted to *Chusquea* bamboo thickets. Its habitat has undergone extensive clearance for pasturelands. The known population is of 24 territories with an estimated extent of occurrence of 5.8 km<sup>2</sup>, giving a cautious global population estimate of 57-156 territories. We propose that the IUCN treat the new species as Critically Endangered. Although protected in the Colibri del Sol reserve, it is in need of further conservation actions.

## Resumen

Una nueva especie de Tororoi *Grallaria* (familia Grallariidae) se describe en la Reserva Natural de las Aves Colibrí del Sol, situada en la vertiente sur-oriental del macizo del Páramo del Sol en el extremo norte de la Cordillera Occidental de Colombia. El Tororoi de Urrao *Grallaria fenwickorum* sp. nov., se parece más a *G. milleri*, la cual es endémica de la Cordillera Central, pero tiene el pecho gris y carece de una banda morena en el pecho como existe en *milleri*. El canto de la nueva especie, se constituye de tres notas, que se incrementan en longitud y frecuencia acústica. El canto posee notas que son más cortas, bajas y de una diferente forma con respecto a *G. milleri*. Adicionalmente, el llamado de la nueva especie es más corto y bajo. *G. fenwickorum* y está restringido a bosque de niebla montano (2.600-3.300 m), en donde sus territorios son restringidos a matorrales de bambú *Chusquea*. Su hábitat ha sido sujeto a deforestación extensiva, especialmente para obtención de pastizales. La población conocida es de 24 territorios con una estimativa del área de ocupación de 5,8 km<sup>2</sup> y una estimativa de población mundial entre 57-156 territorios. Se propone que la IUCN pueda categorizar la nueva especie como en estado Crítico. Aunque la especie se encuentra protegida dentro la reserva Colibrí del Sol, es necesario llevar a cabo adicionales acciones de conservación.

## Introduction

The municipality of Urrao in southwestern Antioquia department includes the 175 km<sup>2</sup> Páramo del Sol massif, also known as Páramo Frontino, that is the highest and largest massif of the 750 km long Western Andes. The massif is a

partially eroded caldera complex with high-level, volcanic-hosted heavy metal mineralization from Upper Tertiary intrusions, including dense high-grade gold, zinc and copper deposits that have attracted the attention of mining companies. The

massif also holds over 2,700 ha of relatively intact páramo and *Polyblepis* woodland that represents an area greater than all the other páramos in the Western Andes, combined.

Despite the unique geological and environmental characteristics and its location 55 km west of Colombia's second largest city, Medellín, the massif only received brief previous ornithological exploration in December 1951 when Melbourne A. Carriker, Jr. visited the páramo and discovered a new hummingbird species later described as Dusky Starfrontlet *Coeligena orina* (Wetmore 1953). For over 50 years thereafter, the site remained unexplored, largely due to security concerns, until Fundación ProAves launched an expedition to survey the massif (Florez *et al.* 2004, Krabbe *et al.* 2006). This expedition resulted in the rediscovery of *Coeligena orina* on 9 August 2004 (Florez *et al.* 2004, Krabbe *et al.* 2005) and the first sound recordings of *Scytalopus canus* (Krabbe & Cadena 2010) as well as a host of range extensions (Krabbe *et al.* 2006).

In addition to scientific findings, it became clear that the Páramo del Sol is a key area for threatened birds requiring protection from the expansion of mining and pasturelands on the massif's mineral-rich and fertile soils. In 2005, thanks to the personal support of the Fenwick family through American Bird Conservancy, Fundación ProAves acquired and protected over 731 has of the montane forest and páramo. This established the Colibrí del Sol Bird Reserve, named after the Critically Endangered Dusky Starfrontlet *Coeligena orina*.

With the continuing support of American Bird Conservancy and USFWS Neotropical Migratory Bird Conservation Act, ProAves implemented a series of research and monitoring stations across the reserve, primarily with banding transects. During banding sessions on February 21 and March 30 2008, ProAves researcher Diego Carantón-Ayala (DCA) and reserve manager Luis

Rubelio García (LRG) caught a mystery *Grallaria* antpitta. This *Grallaria* was also tape-recorded in late 2008 by Alonso Quevedo. The population was considered to be a new species and was discussed as such in the recent description of *Grallaria milleri gilesi* (Salaman *et al.* 2009a). It was added to the Colombian checklist as “*Grallaria sp.*” (Donegan *et al.* 2009, Salaman *et al.* 2009b) and photographs have been published online (e.g. Fundación ProAves 2009). Since 2008 to date, the authors along with many ornithologists and birders have seen, photographed, tape-recorded and studied the distinct new antpitta at Colibrí del Sol Bird Reserve. A family party is seen daily feeding at a worm-feeding station beside the reserve lodge alongside Chestnut-naped Antpitta *G. nuchalis*.

Based on plumage characteristics and differences in vocalizations, the *Grallaria* at Colibrí del Sol Bird reserve represents a new taxon, which meets the requirements for species status under the Biological Species Concept (Johnson *et al.* 1999, Helbig *et al.* 2002), which we hereby name:

### ***Grallaria fenwickorum* sp. nov.**

#### **Fenwick's Antpitta Tororoi de Urrao**

#### **Holotype**

The holotype is constituted solely by:

- a) Feather samples (total of 14 feathers from the wing, tail and body) deposited at the Museo de Historia Natural Jose Celestino Mutis, Facultad de Ciencias de la Universidad de Pamplona, tissue collection No.699 (**Figure 1**).
- b) For purposes of Article 74.1.4 of the International Code for Zoological Nomenclature, to the extent applicable, the individual depicted in Figure 1 and the Cover of this edition of Conservación Colombiana.

These materials are based on an adult tape-recorded, captured and banded (with ProAves ring no. D001108), from which feather samples

were taken, and plumage description was taken using Munsell (1977). These steps were taken in the field and the bird was photographed before being released alive on 11 January 2010 by LFB and LRG. The individual was captured within the Colibrí del Sol Bird Reserve, Vereda El Chuscal, Municipality of Urrao, Department of Antioquia (06°25'53.1"N 76°04'57.9"W). Elevation 3,130 meters asl.

An extensive further series of photos of the individual on which the holotype is based can be downloaded at: <http://www.flickr.com/photos/proaves/sets/72157623898966996/>. Tape-recordings of the individual on which the

holotype is based can be downloaded at: <http://www.xeno-canto.org/48114>

Measurements of individual on which the holotype is based, taken in the field are set out in Table 1. No notable subcutaneous fat in the sternum area and limited body moult were noted.

Although two full specimens of the new species are available at ICN (see “other specimens” below), we do not consider it appropriate to designate these as type specimens given the circumstances in which they were collected (see Comité Editorial de Conservación Colombiana 2010 for further details).

**Figure 1:** Feather sampled from the holotype of Fenwick's Antpitta (*Grallaria fenwickorum*) from the Colibrí del Sol Bird Reserve, Colombia, including a combination of wing, tail and mostly body feathers.



Feather samples constitute a ‘*part of*’ an ‘*example of animal*’ for purposes of the definition of ‘specimen’ in the ICBN Code. The term ‘specimen’, as so defined, is referred to in the

definition of the term ‘*holotype*’ in the ICBN Code. Consequentially, the feather samples taken in this instance constitute a valid holotype on any interpretation of the Code (Donegan 2008b). A

similar procedure of the designation of feather samples as a holotype was recently applied by Athreya (2006).

There have been various recent concerns expressed about the validity of descriptions where the holotype is designated as an individual depicted in a published photograph, as opposed to a specimen or sample (e.g. Timm et al. 2005, Dubois & Nemésio 2007). Although such criticisms have been debunked by members of the Commission's secretariat (e.g. Wakeham-Dawson et al. 2002, Polaszek et al. 2005), we recognize that there are controversies surrounding the use of illustrated individuals as holotypes and our designation of the individual illustrated in Figure 1 as also forming part of the holotype is therefore qualified accordingly.

## Diagnosis

Referred to *Grallaria* Vieillot, 1816 (type: Variegated Antpitta *G. varia*) due to: its medium to large size; long tarsus; culmen very indistinctly (if at all) ridged and gradually but strongly curved from the base; scutellate tarsus; rictal bristles distinct but slender; and chin and upper-throat feathers with long, slender terminal setae (Lowery & O'Neill 1969).

*G. fenwickorum* is typical member of the plain-coloured group (proposed subgenus *Oropeza* Ridgway 1909), due to its relatively small wing (75–103 mm); fairly uniform upperparts and underparts (considered separately) without strong streaks, squamations or bars; tail more than half as long as wing (tail / wing ratio 0.54–0.60); inner edge of tarsus distinctly convolute; and presence of 12 rectrices (Lowery & O'Neill, 1969).

*G. fenwickorum* is considered probably most closely related to Brown-banded Antpitta *G. milleri* due to similarities in voice and biometrics and generally non-descript plumage. There are two described subspecies of *G. milleri*: *G. m. milleri* occurs 165 km southeast in the Central Andes, and the larger *G. m. gilesi* occurs 70 km

east in the northern sector of the Central Andes. *G. fenwickorum* can be easily diagnosed from both taxa by the complete lack of a brown breast band (with the breast instead being uniform slate grey) and lighter brown dorsal plumage. Its vocalizations are also distinct (see below).

Based on published values for *G. milleri* from Salaman et al. (2009a) and Kattan & Beltrán (1999), *G. fenwickorum* appears to average longer wing, tail, bill and tarsus than *G. m. milleri* (Table 1), being more similar to *G. m. gilesi* (as noted in Salaman et al. 2009a). It also appears larger bodied than *G. m. milleri*.

## Description of the holotype

The following is based on Munsell Color (1977) codes. *Face*: Head overall uniform Olive Brown (5Y 5/6), with light gray lores and thin eye-ring (5Y 7/2) with black feather tips. Ear-coverts, malar, and nape warmer cinnamon-brown (10YR 3/6). *Upperparts*: Entire upperparts, from the forecrown to the uppertail-coverts and shoulders, uniform Olive Brown (5Y 5/6). Tail darker brown (7.5YR 4/2). *Wings*: upperwing coverts and flight feathers uniform Olive Brown (7.5YR 7/10) with underwing coverts warm cinnamon-brown (2.5Y 6/4). *Underparts*: Throat light gray (5Y 7/2) with darker olive brown tips to the feathers. Lower underparts and flanks to undertail-coverts pale grey (2.5Y 8/2) with paler gray feather tips and slight olive gray (5Y 6/2) tinge to feathers on the side of the chest and being paler whitish gray on the abdomen. Thighs pale grayish olive-brown (5Y 5/4). *Soft parts* Tarsi and feet bluish-gray. Iris dark brown. Mandible bluish-horn with paler tip and black maxilla.

## Other specimens

There are two specimens of *G. fenwickorum* that were collected without the necessary permit within the Colibrí del Sol Bird Reserve by Diego Carantón-Ayala. There is an on-going investigation by the regional environmental authority as to the incident. These specimens are not designated as type material given the

circumstances in which they were collected and the possibility that they could be confiscated by the authorities or decommissioned from the relevant collection. Although we considered Donegan (2008b)'s recommendation that '*If a specimen based on a dead organism exists, then it is better to use it for a type specimen, whatever the circumstances in which it was procured*', we have decided against such an approach in this instance and instead designated a different holotype for which the relevant permit was available. However, the two specimens mentioned are currently available to be inspected and studied and therefore support the rationale for the description. These specimens are illustrated in **Figure 2**. Details of the specimens are as follows:

1. Adult male (80% skull ossification), catalogue no. 36689, held in the ornithological collection of the Instituto de Ciencias Naturales, Universidad Nacional de Colombia (ICN-MHN), Bogotá. Collected on 30 March 2008 by DCA in the Colibrí del Sol Bird Reserve, Vereda El Chuscal, Municipality of Urrao, Department of Antioquia ( $6^{\circ}25'38.9''\text{N}$ ,  $76^{\circ}05'13.1''\text{W}$ ). Elevation 2,850 meters asl. Well developed testes, no subcutaneous fat. Stomach contents included insects (mainly coleopteran remains).

2. Immature male (30% skull ossification), catalogue no. 36688 held in ICN-MHN. Collected on 20 February 2008 by DCA at the same locality as the adult specimen but 2,820 meters asl. Enlarged testes, no subcutaneous fat. Stomach contents included insect remains.

Should the status of these specimens become resolved, then the authors would encourage, or be willing, to designate the adult male as the neotype. Less stringent conditions should apply to neotype designations where a type specimen other than a full specimen is used (Donegan 2008b).

### Additional individuals

At least 10 territories are known within the Colibrí del Sol Bird Reserve and one territory just outside the reserve in an isolated 1 ha fragment of forest. We obtained many detailed field sightings of *G. fenwickorum*, especially at the worm-feeding station where prolonged views in good light and close quarters are obtained daily. There is no noticeable variation between individuals and pairs, suggesting that the species is sexually monomorphic as in other *Grallaria*.

**Table 1:** Comparison of morphometrics of *G. fenwickorum* with *G. m. gilesi*, *G. m. milleri* (specimens unless otherwise stated) published by Salaman *et al.* (2009a). The mean is given, followed by the standard deviation and then the range, with Kattan & Beltrán (1999) data also presented for comparison. There is expected to be small differences between data as a result of different measuring techniques.

	Flat wing chord (mm)	Tail length (mm)	Tarsus length (mm)	Maxilla (mm)*	Body mass (g)	Tail / tarsus ratio	Wing / tarsus ratio
<i>G. fenwickorum</i> (holotype)	99	57	49.9	22.9	53.5	1.14	1.98
<i>G. fenwickorum</i> (ICN 36689)	95	63	46.5	23.5	56.4	1.35	2.04
<i>G. fenwickorum</i> (ICN 36688)	96	63	44.5	22.4	57.4	1.41	2.15
<i>G. m. gilesi</i> (holotype)	97	55.8	40.8	18.8	N/A	1.37	2.38
<i>G. m. milleri</i> (n=10)	$88.6 \pm 2.9$ (85.0–94.5)	$49.0 \pm 2.9$ (43.6–53.0)	$44.5 \pm 0.6$ (43.4–45.5)	$21.0 \pm 1.9$ (19.3–24.7)	N/A	$1.10 \pm 0.06$ (0.98–1.19)	$1.99 \pm 0.08$ (1.88–2.13)
<i>G. m. milleri</i> (Kattan & Beltrán 1999; n=18 live individuals & specimens)	$89.6 \pm 3.5$	N/A	$46.6 \pm 2.4$	$17.1 \pm 1.5$	52±3.2	N/A	Based on means: 1.92

**Figure 2:** Above *G. m. milleri* ICN 35692 (Caldas, Manizales, Río Blanco, 15/11/2005); Middle: *G. fenwickorum* ICN 36689; Bottom: *G. fenwickorum* ICN 36688.



**Figure 3:** *Grallaria fenwickorum* pullus (c. 18 days old) covered with down on June 12 2008.



One pullus (just fledged) covered with down and only moderate unsheathing of flight feathers was captured, photographed (**Figure 3**) and released on June 12 2008 by DCA and LRG. Additional photos are online at: <http://www.flickr.com/photos/proaves/sets/72157623898966996/>). The mandible and commissures were conspicuous bright orange, while the tarsi and feet were dark pink. The entire upperparts, head and chest were black above with broad rufous-brown tips to give a barred appearance, while the lower underparts (lower breast to undertail coverts) were creamy-buff colored. The flight feathers were partially grown and sheathed.

### **Etymology**

The epithet honors the Fenwick family; George, Rita, Cyrus, Sarah and Rachael Fenwick of The Plains, Virginia, USA. George Fenwick is the President of American Bird Conservancy (ABC)

and Rita Fenwick is the Development Director of ABC – two amazing individuals that have been a force for conservation in the USA and across Latin America, especially Colombia. Importantly, the Fenwick family dedicated their own personal resources to support Fundación ProAves buy and save the 731 ha of Páramo del Sol, establish the Colibrí del Sol Bird Reserve and fund the research which led to this discovery. This commitment to bird conservation has saved many bird species, and now, one more - *Grallaria fenwickorum*. To the Fenwick family – thank you and long live bird conservation. The Spanish name represents the municipality of Urrao where the species may be restricted and where it is hoped that the local community will rally to help save the species. The name *fenwickorum* is not known to be used for any other Grallariidae or suboscine passerine.

### **Notes on authors of the new species for purposes of the Code**

Fundación ProAves de Colombia (commonly known as ‘Fundación ProAves’) funded research and conservation work which led to this description. It is not unusual for an institution to be named as an author of a scientific publication (see e.g. Timm et al. 2005 cited above). However, in this instance, the authors and Fundación ProAves de Colombia have agreed that it would not be appropriate for Fundación ProAves de Colombia to be treated as an author of the name *fenwickorum* for purposes of the Code.

Accordingly, the full citation of the species’ name should be:

*Grallaria fenwickorum* Barrera & Bartels, 2010.

### **ICZN Code of Ethics**

The authors of this article (or their representatives) have communicated with Diego Carantón regularly and on numerous occasions over a period of greater than one year (in this case, starting 19 months before the date of this publication, which the authors consider to be a

more than reasonable period in the circumstances) in relation to the description of the new species of *Grallaria* described herein, and Sr. Carantón has failed to establish a new name for this species within that period. As a result, the authors have complied with the requirements of paragraph 2 of the ICZN Code of Ethics as regards that person.

## Vocalizations

The song and calls of *G. fenwickorum* were recorded at multiple locations in the region of the type locality by the first two authors (n=6 recordings) and compared with those of all other members of the genus *Grallaria* on [www.xenocanto.com](http://www.xenocanto.com). Both the song and call clearly most closely resemble that of *G. milleri* but could not be confused with any other antpitta species. The recordings of *G. fenwickorum* were studied and compared with recordings of *G. milleri* available on [www.xeno-canto.org](http://www.xeno-canto.org), Alvarez et al. (2007) and Isler & Whitney (2002).

The vocal variables set out in Table 2 were studied. Both songs and calls were analyzed. Only the first three songs or calls for each available recording were studied. Data on the acoustic variables studied is set out in Table 2.

The song of *Grallaria fenwickorum* consists of three notes, progressively increasing in acoustic frequency and length, with the first note briefest and lowest, and the final note longest and highest. The song could be transcribed as: “tu, tuut, TUUET”.

The song of *G. milleri* is similar to that of *G. fenwickorum*, also comprising three notes of progressively higher pitch and greater length (**Figure 4**). However, each note of the *G. milleri* song is diagnosably longer and higher pitched than the equivalent note in *G. fenwickorum*. There are average differences in the length of intervals between notes (shorter in *milleri*) and total song duration (shorter in *milleri*) but songs are not fully diagnosable on these variables. There are also differences in note shape, with notes in *G. fenwickorum* songs appearing more as an up-down stroke and with longer “tails” compared with a flatter, more varying note shape in *G. milleri*.


We treat the acoustic frequency variables for each of the three notes as correlated; and the note duration of each of the three notes also has being correlated. As a result, these differences constitute two vocal differences (one note length based, one frequency based) for purposes of assessing species limits, with differences in note shape constituting a third difference. The diagnostic differences in acoustic frequency, note duration and note shape between *G. milleri* and *G. fenwickorum* therefore meet the suggested benchmark for species limits in Grallariidae (Donegan 2008a).

The call of *G. fenwickorum* is a sharp, loud, single note which is higher-pitched than the song (**Figure 5a**). The call is frequently heard in response to imitation of the song and playback. The call is similar to that of *G. milleri*, but differs in being shorter and lower-pitched with a less curved note shape (**Figure 5b**).


**Table 2:** Vocal variables of *G. m. milleri* and *G. fenwickorum* songs and calls. Data is presented in the form Mean  $\pm$  standard deviation (lower recorded limit – upper recorded limit) (n=sample number). Diagnosability is based on Donegan (2008)'s “Level 5” test (Isler et al. 1999's diagnosability test).

Songs	<i>G. milleri</i>	<i>G. fenwickorum</i>	Diagnosable?
Number of notes in song;	3 (all)	3 (all)	No
Total song duration (s)	1.02 $\pm$ 0.11 (0.85-1.14) (n=10)	0.83 $\pm$ 0.02 (0.80-0.85) (n=16)	No
Song speed (No. of notes /s).	2.98 $\pm$ 0.33 (2.63-3.52) (n=10)	3.62 $\pm$ 0.07 (3.54-3.77) (n=16)	No
Duration of first note (s)	0.21 $\pm$ 0.02 (0.19-0.24) (n=10)	0.10 $\pm$ 0.01 (0.07-0.11) (n=15)	Yes
Duration of second note (s)	0.22 $\pm$ 0.02 (0.19-0.25) (n=10)	0.10 $\pm$ 0.01 (0.09-0.12) (n=17)	Yes
Duration of third note (s)	0.22 $\pm$ 0.02 (0.17-0.25) (n=10)	0.12 $\pm$ 0.01 (0.10-0.14) (n=17)	Yes
Duration of interval between first and second note (s)	0.28 $\pm$ 0.11 (0.14-0.43) (n=10)	0.37 $\pm$ 0.01 (0.35-0.41) (n=16)	No
Duration of interval between second and third note (s)	0.09 $\pm$ 0.03 (0.06-0.14) (n=10)	0.12 $\pm$ 0.01 (0.11-0.15) (n=17)	No
Acoustic freq. of 1 <sup>st</sup> note (kHz)	2.65 $\pm$ 0.08 (2.55-2.77) (n=10)	2.33 $\pm$ 0.03 (2.30-2.38) (n=16)	Yes
Acoustic freq. of 2 <sup>nd</sup> note (kHz)	2.80 $\pm$ 0.07 (2.69-2.91) (n=10)	2.47 $\pm$ 0.05 (2.40-2.60) (n=17)	Yes
Acoustic freq. of 3 <sup>rd</sup> note (kHz)	3.02 $\pm$ 0.05 (2.91-3.08) (n=10)	2.76 $\pm$ 0.04 (2.69-2.86) (n=17)	Yes
Calls			
Call duration (s)	0.45 $\pm$ 0.01 (0.41-0.47) (n=18)	0.32 $\pm$ 0.01 (0.30-0.36) (n=19)	Yes
Minimum acoustic freq. (kHz)	4.62 $\pm$ 0.15 (4.32-4.75) (n=18)	4.25 $\pm$ 0.04 (4.18-4.35) (n=19)	No
maximum acoustic freq. (kHz)	5.46 $\pm$ 0.21 (5.10-5.78) (n=18)	4.90 $\pm$ 0.06 (4.77-5.00) (n=19)	Yes
variation in freq. (kHz)	0.84 $\pm$ 0.09 (0.72-1.05) (n=18)	0.65 $\pm$ 0.06 (0.54-0.73) (n=19)	No

**Figure 4:** Territorial song of the holotype of *G. fenwickorum* (b), recorded by LFB on 11 January 2010 in Colibrí del Sol Bird Reserve and (a) *G. m. milleri* recorded by Nick Athanas on 20 February 2007 in El Mirador Bird Reserve, Quindío Dept (04°08'22"N 75°44'08"W), No.:XC10721. Accessible at [www.xeno-canto.org/10721](http://www.xeno-canto.org/10721)



**Figure 5:** Calls of (a) *G. fenwickorum* recorded by LFB on 11 January 2010 in Colibrí del Sol Bird Reserve and (b) *G. m. milleri* recorded by Frank Lambert on 29 may 2007 in El Mirador Bird Reserve, Quindío Dept, No.:XC16777. Accessible at <http://www.xeno-canto.org/recording.php?XC=16777>



## Distribution

*Grallaria fenwickorum* is known only from the immediate vicinity of the type locality of Colibrí del Sol Bird Reserve (**Figure 6**), on the southeastern slope of the Páramo del Sol massif, in the Municipality of Urrao, Department of Antioquia (06°25'N 76°04'W). Here, the Western Andes reaches its highest elevations. Other high elevation species, such as *Scytalopus canus* and *Coeligena orina* appear also to be restricted to the northern massifs of the western Andes and do not extend in distribution south to other high massifs in the range, such as Cerro Munchique in Cauca, which has been subject to extensive ornithological efforts.

We have documented eleven territories, of which only one has been found outside of the reserve (620 meters to the southeast in a isolated 1 ha forest patch), but all on the slightly less humid


eastern slope of the Western Andes. *G. fenwickorum* appears restricted to montane cloud forest dominated with *Quercus humboldtii* (Colombian Oak) between 2,600-3,300 meters elevation where territories are largely restricted to *Chusquea* bamboo thickets. It is interesting to note that the local district where the type-locality and reserve occurs is called “El Chuscal”, referring to the extent of *Chusquea* bamboo!

We suspect the range of *G. fenwickorum* to be larger than just the Colibri del Sol reserve, but surveys 45 km to the south on the eastern slope of the West Andes within its elevational range failed to locate the species. Furthermore, there have been many surveys of the montane forests of the West Andes by ornithologists and birders in the past decade, all of which have failed to report the species (Cuervo et al. 2003). This would suggest that *G. fenwickorum* has a particularly restricted

range, probably based on its unique ecological preferences on the more sheltered and slightly less humid eastern slopes of the West Andes, an area that has been severely impacted by deforestation. We consider that the species could range lower in elevation, below 2,500 m, however there is virtually no surviving humid montane forest below that elevation on the eastern slope of the West Andes, so the hopes of a viable population at lower elevations looks unlikely.


**Figure 6.** Distribution map of *G. fenwickorum* in (a) Google Maps terrain image of the northern sector of the Western and Central Andes and (b) LandSat image of the Páramo del Sol massif in the Municipality of Urrao, Dept. of Antioquia, Colombia with territories shown (yellow spots).

A)



We hope that additional populations survive, with an important site for further studies being the eastern slopes of Farallones de Citará. However, this peak is highly exposed and much more humid. We hope that the availability of sound recordings of the new species in [www.xenocanto.com](http://www.xenocanto.com) will aid birders' and ornithologists' searches for the species and hopefully result in the discovery of new populations.

B)



### Taxonomic affinities

The recent description of *G. milleri gilesi* and discussion of morphological features within *G. milleri* (Salaman *et al.* 2009a) allows light to be shed on the taxonomic affinities of *G. fenwickorum*. Based on morphological and vocalization characteristics described above, the closest relative of *G. fenwickorum* is clearly *G. milleri*. The new species is most probably most closely related to *G. m. gilesi*, which is larger in biometrics, being more similar to *G. fenwickorum* in this respect. Furthermore, *G. m. gilesi* is suspected formally to have occurred just 70 km east on the western slope of the northern sector of the Central Andes, whilst *G. m. milleri* occurs

further away - 165 km southeast in the middle sector of the Central Andes. There are no available vocalizations of the possibly extinct subspecies *G. m. gilesi* (Salaman *et al.* 2009).

### Population and territory size

Surveys were conducted, consisting of transect point counts, using playback, between 2,780 m (the lowest of contiguous forest) and 3,345 m. The area surveyed was approximately 98 ha within the core area of known territories within the reserve. Most points were at intervals of at least 50 m apart, the exceptions occurring in habitat not considered to be prime (Type A; see Habitat details below) where no response to

playback was solicited. No two points that received responses were closer than 50 meters. In a few cases adjacent points received responses from the same area, in which case this was counted as one territory. In total 85 separate points were surveyed over 14 days between January 11 and February 2, 2010. All point counts were conducted between 0615-1030 hrs, though rarely after 0945hrs, and 1600-1730 hrs. The maximum period of playback at a point was ten minutes, roughly following a 30 seconds on, 30 seconds off pattern, though later on the usage of playback lessened, with more time between each playing as the birds often responded 1-3 minutes after playback stopped. Both song (60%) and call (40%) were used.

The transect point counts identified approximately 24 territories of *G. fenwickorum*. Due to time restrictions, an accurate size of each territory was not obtained, but based on the proximity of territories and micro-habitat boundaries it is estimated that the average territory area between 0.4-1.1 ha, which is consistent for territory size of *G. milleri* of an estimated 0.5-5.4 ha (Kattan & Beltran 2002). It is also estimated that the total area of occupancy was between 9.6 to 26.4 ha surveyed with an occupancy of 9.8-26.9% of the area surveyed.

Extrapolating habitat preferences across the reserve and adjacent suitable habitat based on contour maps of the eastern and southeastern slope of Páramo del Sol massif, *G. fenwickorum*'s estimated extent of occurrence is 5.8 km<sup>2</sup>, giving a cautious estimate of between 57-156 territories. Each territory refers to a singing adult male, so does not necessarily translate to an estimate of the number of pairs.

## Breeding

Territorial vocalizations appear more pronounced in the first half of the year, while evidence of breeding can only be confirmed between May-June, with one recently fledged pullus (estimated to be c. day 18), covered with down and only

moderate flight feather unsheathing, was found on June 12 2008. This record coincides with one of the wet seasons, May through June.

## Habitat preferences

*Grallaria fenwickorum* is restricted in a thin altitudinal belt of humid cloud forest dominated with *Quercus humboldtii* (Colombian Oak) between 2600 m and 3300 m, and strongly dependant on *Chusquea* cf *scandens* (Andean Dwarf Bamboo) thickets under the forest canopy or treefall/landslide gaps. The climate at the reserve headquarters (in the core *G. fenwickorum* population area) averages 2,284 mm per annum with a minimum temperature average of 8.8°C and maximum temperature of 16.0°C with the driest months being December-March and wettest from May-June and October-November (unpublished weather station data).

The preferred habitat for *G. fenwickorum* can be classified into three classes based on the extent of *Chusquea* within an area: >90% = Type A, 60-90% = Type B to <20% = Type C habitats. Nineteen territories were found in Type A habitat, three were found in Type B and two in Type C. As the area studied is very mountainous, virtually all territories were on slopes averaging 20-30% grade. However, a few territories contained areas up to 60% grade while others near Quebrada Santa Barbara included areas with 0-5% grade. Photos of the various habitat types at are shown online at: <http://www.flickr.com/photos/proaves/sets/72157623898966996/>.

**Type A (nineteen territories):** the main habitat type is considered to be of early successional vegetation from large treefall sites or landslides and consists of thickets of dense, mature *Chusquea* bamboo with low light penetration. The low lighting is not generally due to a dense canopy but rather a fairly sparse upperstorey (trees sometimes spaced as far apart as 15-20 m) allowing *Chusquea* to flourish and dominate, creating a dense to very dense understorey "canopy" at 1.5-3 m above the forest floor. A

dominant characteristic of this habitat is the presence of clusters of horizontal to near horizontal dead and living *Chusquea* stalks with accumulated dead foliage that *G. fenwickorum* favors for cover. The relatively open forest floor, that most individuals preferred, was generally covered in a thin layer of dead *Chusquea* leaves, at most 5cm thick but usually averaging about 2cm. Height of the upperstorey (loosely equivalent to canopy) varied from 7-25 m and did not appear to be a factor influencing the density or location of territories.

**Type B (three territories):** very similar to Type A habitat with dense patches of *Chusquea*, but mixed patchily with other plant species, such as ferns and very young trees. Type B is typically representative of advancing successional vegetation.

**Type C (two territories):** two territories occurring along the same stream between 2815 and 2860m with very different habitat from Types A and B. The canopy is denser and lower, between 7-12m, also with a very dense understorey, between 0-1.5m, dominated by ferns (30% at one territory, 60% at other) and herbaceous plants with a limited presence (<5%) of *Chusquea* in the immediate area, although both territories had stands of *Chusquea* nearby (20m and 30m, respectively), that the individuals may or may not have been using. The forest floor was covered in a fairly thick layer of dead foliage, averaging between 5cm and 10cm.

## Behavior

*Grallaria fenwickorum* exhibits typical behavior as seen by other members of the genus, being a shy terrestrial antpitta that forages on insects in the leaf-litter and dense understorey and ascends to perches up to 1.5 meters off the ground to sing from within the dense bamboo understory. The species is most active and vocal from dawn (~6.00 hrs) to 9.45 hrs and again from 16.00 hrs to dusk (~18.30 hrs).

The species responds well to playback throughout the year, however, individuals often hard to see. In order to minimize disturbing this very shy species with playback for visiting birders, LRG established a worm feeding station that immediately proved very successful. At mid-morning each day, worms are placed on two trays and birds are called up. Individuals, pairs and occasionally 3-4 individuals (assumed to be young) bound down the slopes to feed on the worms. *G. fenwickorum* territories occasionally overlap with the much larger Chestnut-naped Antpitta *G. nuchalis* (a species which is at least twice as large) which also frequents the worm-feeding station.

## Threats

*Grallaria fenwickorum* is presently known from 24 territories with an estimated extent of occurrence is 5.8 km<sup>2</sup>, giving a cautious estimate of between 57-156 territories, although it is unclear if all those territories are breeding pairs. The known and estimated population is extremely small and of great conservation concern. Furthermore, *G. fenwickorum* has highly specific ecological requirements, being dense *Chusquea* thickets within primary montane cloud forest on less humid east-facing or sheltered valleys of the Western Andes. Very few peaks of the height of Páramo del Sol exist and most of those are considerably further south in the West Andes (e.g. Munchique, Farallones de Cali). Those peaks have been surveyed extensively with no records of this new species. As a result, the species is likely restricted to high elevation peaks in the northern part of the West Andes only.

The fertile eastern slopes and valleys of the Western Andes have long been extensively deforested for rich pasture and agricultural lands primarily as it is less humid than the adjacent western or Pacific slope of the Western Andes. Even steep slopes have been cleared in the region, assisted by burning forests upslope during extreme droughts (typically El Niño years). The extent of montane cloud forest that survives

between 2,000-3,300 meters on the eastern slope of the Western Andes in the department of Antioquia is estimated at less than 21% of its original cover (approximately 978 km<sup>2</sup>). However, much of this is heavily fragmented and less than 0.16% is protected.

One core area of surviving humid montane forest dominated by *Quercus humboldtii* is the eastern slope of the Páramo del Sol massif, particularly the upper Río Urrao watershed, including Quebrada Santa Barbara that runs through the Colibrí del Sol Bird Reserve. The Río Urrao watershed totals 6,050 ha of which 37% is primary montane forest with moderate levels of forest fragmentation and of the surviving forest, the Colibrí del Sol Bird Reserve protects 19.2% (428 ha of forest). Examination of forest cover change between 2000 and 2007 within the Urrao watershed documented a loss of 83 ha of forest or 3.6% of forest cover, despite a gain in reforestation of 38 ha of pasturelands to forest in the Colibrí del Sol Bird Reserve.

Unfortunately, the partially eroded caldera complex underlying Páramo del Sol features high-level, volcanic-hosted heavy metal mineralization from Upper Tertiary intrusions (Aspden *et al.* 1987, MiningLife 2004). A significant improvement in public order within the region in recent years together with documented high grade gold, zinc and copper deposits hosted within the massif have attracted the attention of several mining companies. In August 2007, Continental Gold de Colombia S.A., a Canadian subsidiary of Continental Gold Ltd., secured mineral exploration rights to 500 ha area within the Colibrí del Sol Bird Reserve. This exploration was successfully disputed, but given that the entire Páramo del Sol massif is privately owned, land speculation has been rife. The likelihood of protecting additional forests in the area faces significant challenges.

## Proposed IUCN status

Based on the information published within this article, *G. fenwickorum* meets the IUCN criteria:

### **CRITICALLY ENDANGERED** B1a+b(i,iii,v); C2a(ii):

--B. Geographic range in the form of B1 (extent of occurrence) estimated to be less than 100 km<sup>2</sup>, and (a.) known to exist at only a single location and (b.) projected decline in (i) extent of occurrence, (iii) area, extent and/or quality of habitat, and (v) number of mature individuals, and

--C. Population size estimated to number fewer than 250 mature individuals and (2.) a continuing projected decline in numbers of mature individuals (a.) Population structure in the form of (ii) at least 90% of mature individuals in one subpopulation.

It is noteworthy that the closest relative of *G. fenwickorum*, *G. milleri* is considered Endangered (EN: B1a+b(iii,v), VU: C2a(i); D1) with an estimated extent of occurrence of 660 km<sup>2</sup> and known from ten locations across the Central Andes (Salaman *et al.* 2009, BirdLife International 2010), while *G. m. gilesi* is considered probably Extinct (Salaman *et al.* 2009). This indicates that this group of antpittas are particularly sensitive to environmental conditions and prone to extinction.

## Conservation Actions

While *Grallaria fenwickorum* is effectively protected within the 731 ha Colibrí del Sol Bird Reserve that is owned and managed by Fundación ProAves and registered within the National System of Protected Areas, the species requires a significantly larger area of suitable habitat protected while suitable forests still survive. It is also essential that searches should be made for *G. fenwickorum* along the eastern slope of the Western Andes to locate possible new populations.

Considering the critical importance of protecting *G. fenwickorum* and many other threatened and endemic species, American Bird Conservancy and Fundación ProAves will shortly publish a comprehensive Conservation Action Plan for the Páramo del Sol massif, based on extensive biodiversity surveys to identify key biodiversity sites. This plan will lay a strong framework for future actions, using the toolbox of conservation actions to ensure the strategic and effective protection of *G. fenwickorum* as well as many other threatened birds (e.g. Dusky Starfrontlet *Coeligena orina* [CR], Paramillo Tapaculo *Scytalopus canus* [proposed CR], and Chestnut-bellied Flowerpiecer *Diglossa glorriasisima* [EN]) and amphibians (e.g. Nicefori's Harlequin Frog *Atelopus nicefori* [CR], *Bolitoglossa medemi* [VU], *Hyloxalus borjai* [DD]) that depend on this critical Alliance for Zero Extinction site.

## Acknowledgements

We acknowledge Diego Carantón and Luis Rubelio García, who first collected the new species whilst working for ProAves. The authors are grateful for the support of many people that made this article possible, including Alonso Quevedo, David Caro, Carlos Andres Páez, Nick Athanas, Sara Inés Lara and the other sound recordists and observers that helped the authors in the field. Thanks to Alonso Quevedo, Thomas Donegan, and David Caro for their comments on the paper; and to the editorial committee of Conservación Colombiana for clarifying the circumstances of this description in their editorial. We received research permits from CORPOURUBA (Resolution No. 1043 of 15 July 2006). Paul Salaman, Robert Prys-Jones and Thomas Donegan kindly made available data and photographs used in connection with their studies published in Salaman et al. (2009a) and files for statistical analysis of vocal variables used by Donegan (2008a). We are grateful for the support by Fundación ProAves, American Bird Conservancy, the Fenwick family and USFWS Neotropical Migratory Bird Conservation Act to undertake fieldwork and conservation actions to

protect *Grallaria fenwickorum*. The article was kindly peer-reviewed by David Agro, Paul Salaman and Edward Dickinson, for whose comments we are grateful.

## References

- Álvarez, M., Caro, V., Laverde, O. & Cuervo, A. M. (2007) *Guía sonora de los Andes colombianos*. Instituto Alexander von Humboldt & Cornell Lab. of Orn.
- Aspden, J. A., McCourt, J. M. & Brook, M. (1987) Geometric control of subduction-related magnetism: the Mesozoic and Cenozoic plutonic history of western Colombia. *Journal of the Geological Society* 144: 893-905.
- Athreya, R. (2006) A new species of *Liocichla* (Aves: Timaliidae) from Eaglenest Wildlife Sanctuary, Arunachal Pradesh, India. *Indian Birds*, 2(4), 82-94.
- BirdLife International (2010) Species factsheet: *Grallaria milleri*. Downloaded from <http://www.birdlife.org> on 07/01/2010
- Comité Editorial de Conservación Colombiana (2010) Editorial. *Conservación Colombiana* 12: 4-6.
- Cuervo, A. M., Stiles, F. G., Cadena, C. D., Toro, J. L. & Londoño, G. A. (2003) New and noteworthy bird records from the northern sector of the Western Andes of Colombia. *Bulletin of the British Ornithologists' Club* 123:7-24.
- Donegan, T.M. (2008a) Geographical variation in Slate-crowned Antpitta *Grallaricula nana*, with descriptions of two subspecies, from Colombia and Venezuela. *Bulletin of the British Ornithologists' Club* 128(3): 150-178.
- Donegan T.M. (2008b) New species and subspecies descriptions do not and should not always require a dead type specimen. *Zootaxa* 1761: 37-48.
- Donegan TM. (2009) Type specimens, samples of live individuals and the Galapagos Pink Land Iguana. *Zootaxa* 2201: 12-20.
- Donegan T.M., Salaman P. & Caro, D. 2009. Revision of the status of various bird species occurring or reported in Colombia. *Conservación Colombiana* 8: 80-86.
- Dubois, A. & Nemésio, A. (2007) Does nomenclatural availability of nomina of new species or subspecies require the deposition of vouchers in collections? *Zootaxa*, 1409, 1-22.
- Flórez, P., Krabbe, N., Castaño, J., Suárez, G., & Arango, J. D. (2004) Evaluación avifauna del

- Paramo de Frontino, Antioquia, August 2004. Colombian EBA Project Report Series No. 6, Fundación ProAves, Bogotá, Colombia.
- Fundación ProAves (2009) Récord de observación de aves endémicas en Colombia. *Aleteo* 65.
- Helbig A. J., Knox, A. G., Parkin, D. T., Sangster, G. & Collinson, M. (2002) Guidelines for assigning species rank. *Ibis* 144: 518–525.
- Isler, P. R. & Whitney, B. M. (2002) *Songs of the antbirds*. CDs. Cornell Lab. of Orn., Ithaca, NY.
- Isler M. L., Isler, P. R. & Whitney, B. M. (1999) Species limits in antbirds. The *Myrmotherula surinamensis* complex. *Wilson Bull.* 116: 83–96.
- Johnson, N.K., Remsen, J.V. Jr, & Cicero, C. (1999) Resolution of the debate over species concepts in ornithology: a new comprehensive biologic species concept. Proceedings of the 22 International Ornithological Congress, Durban, pp. 1470–1482.
- Kattan, G. H. & Beltrán, W. (1999) Altitudinal distribution, habitat use, and abundance of *Grallaria* antpittas in the Central Andes of Colombia. *Bird Conservation International* 9: 271–281.
- Kattan, G. H. & Beltrán, W. (2002) Rarity in antpittas: territory size and population density of five *Grallaria* spp. in a regenerating habitat mosaic in the Andes of Colombia. *Bird Conservation International* 12: 231–240.
- Krabbe, N. & Cadena, C. D. (2010) A taxonomic revision of the Páramo Tapaculo *Scytalopus canus* Chapman (Aves: Rhinocryptidae), with description of a new subspecies from Ecuador and Peru. *Zootaxa* 2354: 56–66.
- Krabbe, N., Florez, P. Suarez, G., Castano, J., Arango, J.D., Pulgarin, P.C., Munera, W.A., Stiles, F. G. & Salaman, P. (2005). Rediscovery of the Dusky Starfrontlet *Coeligena orina*, with a description of the adult plumages and a reassessment of its taxonomic status. *Ornitología Colombiana* 3:28-35.
- Krabbe, N., Flórez, P., Suárez, G., Castaño, J., Arango, J. D. & Duque, A. (2006) The birds of Páramo Frontino, West Andes of Colombia. *Ornitología Colombiana* 4: 39–50.
- Lowery, G. H. J. & O'Neill, J. P. (1969) A new species of *Grallaria* from Peru, and a revision of the subfamily Grallarinae. *Auk* 86:1-12.
- Mininglife. (2004) Lunareja Mineral Property on MiningLife. <http://www.mininglife.com>
- Munsell Color. 1977. *Munsell color charts for plants tissues*. 2nd ed. Munsell Color (Firm), Baltimore, MD.
- Polaszek, A., Grubb, P., Groves, C., Ehhardt, C.L. & Butynski, T.M. (2005) What constitutes a proper description: response. *Science*, 309, 2164–2166.
- Salaman, P., Donegan, T. M., & Prŷs-Jones, R. (2009a) A new subspecies of Brown-banded Antpitta *Grallaria milleri* from Antioquia, Colombia. *Bulletin of the British Ornithologists' Club* 129:5-16.
- Salaman P, Donegan T & Caro D. (2009b) Listado de Aves de Colombiana 2009. *Conservación Colombiana* 8: 1-89.
- Timm, R.M., Ramey R.R. II & Nomenclatural Committee of the American Society of Mammalogists (2005) What constitutes a proper description? *Science*, 309, 2163–2164.
- Wakeham-Dawson, A., Morris, S., Tubbs, P., Dalebout, M.L. & Baker, C.S. (2002) Type specimens: dead or alive? *Bulletin of Zoological Nomenclature*, 59(4), 282–286.
- Wetmore, A. (1953) Further additions to the birds of Panama and Colombia. *Smithsonian Miscellaneous Collections* 122: 1-12.

